

Research Article:

**PROTEȚIA NATURALĂ A LEMNULUI
CU CIUPERCI ANTAGONICE**

**NATURAL PROTECTION OF WOOD
WITH ANTAGONISM FUNGI**

Alba ZAREMSKI

Dr.eng. - UR 39 «génétique forestière», CIRAD BIOS TA A-39/C Campus International de Baillarguet
Adresa/Address: F-34398 Montpellier Cedex 5, France
E-mail: alba.zaremski@cirad.fr

Louis GASTONGUAY

Researcher - Institut de Recherche d'Hydro-Québec, Sciences des matériaux
Adresa/Address: 1800 Lionel-Boulet, Varennes, Québec, Canada, J3X 1S1

Yves PRIN

Researcher - Laboratoire des Symbioses Tropicales et Méditerranéennes
Adresa/Address: UMR 113 (CIRAD/INRA/IRD/Agro-M/UM2) TA 10/J, F-34398 Montpellier Cedex 5, France

BIBLIOGRAFIE / REFERENCES

- ALABOUVETTE, C., COUTEAUDIER, Y., LOUVET, J. (1983). Importance des phénomènes de compétition nutritive dans l'antagonisme entre microorganismes. Colloque de la société Française de phytopathologie, Bordeaux (France), 34: 7-16.
- BAKER, R. (1991). Diversity in biological control. *Crop Protection*, 10: 85-94.
- BESSELAT, B. (1985). Résultats obtenus par le service de la protection des végétaux dans le cadre de la lutte contre la pourriture grise de la vigne avec utilisation du *Trichoderma*. L'emploi des ennemis naturels dans la protection des cultures, p. 51-58. INRA, Paris (France).
- BRUCE, A., KING, B. (1983). Biological control of wood decay by *Lentinus lepideus* (Fr.) produced by *Scytalidium* and *Trichoderma* residues. *Material u. Organismen* 18: 171-181.
- BRUCE, A., HIGHLEY, T.L. (1991). Control of growth of wood decay Basidiomycetes by *Trichoderma* spp. and other potentially antagonistic fungi. *Forest Products Journal*, 41(2): 63-57.
- BUTT, T.M., JACKSON, C., MAGAN, N. (2001). *Fungi as Biocontrol Agents: Progress, Problem and Potential*. CABI Publishing, Wallingford, Oxon, UK. p. 398.
- CAVALCANTE, M.S., EATON, R.A. (1981). Inhibition of wood-inhabiting fungi by actinomycetes. International Research Group on Wood Protection, Document No. IRG/WP/1137.
- CHET, I. (1984). Application of *Trichoderma* as a bio control agent. In Proceedings of the 6th congrès Phytopathologie Méditerranéen, Cairo (Egypt), p. 110-111.
- CLAUSEN, C.A., YANG, V. (2007). Protecting wood from mould, decay, and termites with multi-component biocide systems. *International Biodeterioration and Biodegradation*, 59(1): 20-24.
- CLUBBE, C.P. (1980). The colonisation and succession of fungi in wood. International Research Group on Wood Protection: Document No. IRG/WP 1107.
- COMPORATA, M. (1985). Antagonisme in vitro de *Trichoderma* spp vis à vis de *Rhizoctinia solani*, INRA Paris (France), p. 613-620.
- CORBAZ, R. (1990). *Principes de phytopathologie et de lutte contre les maladies des plantes*. Presses polytechniques universitaires. Paris (France), p. 650.
- DAVET, P. (1996). Vie microbienne du sol et production végétale, INRA, Paris (France), p. 383.
- DAVET, P. (1983a). Les *Trichoderma*, exemple de champignon antagoniste d'agents pathogènes. Faune et flore auxiliaire en agriculture. ACTA, INRA- ENSAM- Montpellier (France), p. 193-204.
- DAVET, P. (1983b). Introduction et conservation de *Trichoderma* dans le sol. Les antagonistes microbiens, mode d'action et application à la lutte biologique contre les maladies des plantes. ACTA, INRA- ENSAM- Montpellier (France), p. 159-168.
- DENNIS, C., WEBSTER, J. (1971a). Antagonistic properties of species-groups of *Trichoderma* I. Production of non-volatile antibiotics. *Transactions of the British Mycological Society*, 57(1): 25-39.
- DENNIS, C., WEBSTER, J. (1971b). Antagonistic properties of species-groups of *Trichoderma* II. Production of non-volatile antibiotics. *Transactions of the British Mycological Society*, 57(1): 41-48.

DENNIS, C., WEBSTER, J. (1971c). Antagonistic properties of species-groups of *Trichoderma* III. Hyphal interaction. *Transactions of the British Mycological Society*, 57: 363–369.

DUBOS, B. (1985). L'utilisation des *Trichoderma* comme agent de lutte biologique à l'égard de deux parasites aériens: *Chondrostereum purpureum* (Pers. Ex. fr.) pouzar (plomb des arbres fruitiers) et *Botrytis cinerea* pers. (pourriture grise de la vigne. L'emploi des ennemis naturels dans la protection des cultures, INRA, Paris (France), p. 35-49.

DIK, A.J., ELAD, Y. (1999). Comparison of antagonists of *Botrytis cinerea* in greenhouse grown cucumber and tomato under different climatic conditions. *European Journal of Plant Pathology*, 105: 123–137.

EVANS, P. (2003). Emerging technologies in wood protection. *Forest Products Journal*, 53(1): 14-22.

FREEMAN, M.H., SHUPE, T.F., VLOSKY, R.P., BARNES, H.M. (2003). Past, present, and future of the wood preservation industry. *Forest Products Journal*, 53(10): 8-15.

FREITAG, M., MORRELL, J.J., BRUCE, A. (1991). Biological protection of wood: Status and prospects. *Biodeterioration Abstracts*, 5: 1-13.

HARMAN, G.E., STASZ, T.E. (1989). Combining effective strains of *Trichoderma harzianum* and solid matrix priming to provide improved biological seed treatment systems. *Plant Disease*, 72: 631-637.

HAYES, C.K., PETERBAUER, C., TRONSMO, A., KLEMSDAL, S., HARMAN, G.E. (1993). Antifungal chitinolytic enzymes from *Trichoderma harzianum* and *Gliocladium virens* purification, characterization, biological activity and molecular cloning. In Muzzarelli, R.A.A. (ed). *Chitin Enzymology*. Ancona, Italy, p.383-392.

HJELJORD, L.G., STENSVAND, A., TRONSMO, A. (2000). Effect of temperature and nutrient stress on the capacity of commercial *Trichoderma* products to control *Botrytis cinerea* and *Mucor piriformis* in greenhouse strawberries. *Biological Control*, 19: 149–160.

HIGHLEY, T.L., RICARD, J.L. (1988). Antagonism of *Trichoderma* spp. and *Gliocladium virens* against wood decay fungi. *Material und Organism*, 23(3): 157-169.

HIRATSUKA, Y., CHAKRAVARTY, P., MIAO, S., AYER, W.A. (1994). Potential for biological protection against blue stain in *Populus tremuloides* with a hyphomycetous fungus, *Stachybotrys cylindrospora*. *Canadian Journal Forest Research*, 24: 174-179.

IDWAN SUDIRMAN, L., IRAQI HOUSSEINI, A.I., LE FEBVRE, G., KIFFER, E., BOTTON, B. (1992). Screening of some basidiomycetes for biocontrol of *Rigidoporus lignosus*, a parasite of the rubber tree *Hevea brasiliensis*. *Mycological Research*, 96(8): 621-625.

JIN, X., HAYES, C.K., HARMAN, G.E. (1992). Principles in the development of biological control systems employing *Trichoderma* species against soil-borne plant pathogenic fungi. In Leatham, G.F. (ed): *Symposium on Industrial Mycology*, Mycological Soc. Am., Brock/Springer Series in Contemporary Bioscience, p. 174-195.

LAND, C.J., HULT, K. (1987). Mycotoxin production by some wood-associated *Penicillium* spp. *Letters in Applied Microbiology*, 4: 41-44.

LANUSSE, M., LUNG-ESCAVANT, B., DUBOT, B., TARIS, B. (1983). Etude in-vitro des propriétés antagonistes de 8 espèces de *Trichoderma* à l'égard de deux souches d'*Armillaria mellea*. XXIV colloque de la société Française de phytopathologie, Bordeaux (France), 34: 179-192.

MORRIS, P.I., SUMMERS, N.A., DICKINSON, D.J. (1986). The leachability and specificity of the biological protection of timber using *Scytalidium* sp. and *Trichoderma* spp. International Research Group on Wood Preservation, Document No. IRG/WP/1302.

SHAKERI, J., FOSTER, H.A. (2007). Proteolytic activity and antibiotic production by *Trichoderma harzianum* in relation to pathogenicity to insects, *Enzyme and Microbial Technology*, 40: 961–968.

SMITH, V.L., WILCOX, W.F., HARMAN, G.E. (1990). Potential for biological control of *Phytophthora* root and crown rots of apple by *Trichoderma* and *Gliocladium* spp. *Phytopathology*, 70: 880-885.

STASZ, T.E., HARMAN, G.E., WEEDEN, N.F. (1988). Protoplast preparation and fusion in two biocontrol strains of *Trichoderma harzianum*. *Mycologia*, 80: 141-150.

STILWELL, M.A., WALL, R.E., STRUNZ, G.M. (1973). Production, isolation and antifungal activity of scytalidin, a metabolite of *Scytalidium* sp. *Canadian Journal Microbiology*, 19: 597-602.

VAN DER STEEN, J.J.M., LANGERAK, C.J., VAN TONGEREN, C.A.M., DIK, A.J. (2003). Aspects of the use of honeybees and bumblebees as vector of antagonistic micro-organisms in plant disease control. In: J. Bruin, Editor, *Proceedings of the Netherlands Entomological Society Meeting*, Uitgeverij Nederlandse Entomologische Vereniging. Amsterdam. The Netherlands.

VANNESTE, J.L., HILL, R.A., KAY, S.J., FARRELL, R.L., HOLLAND, P.T. (2002). Biological control of sapstain fungi with natural products and biological control agents: A review of the work carried out in New Zealand. *Mycological Research*, 106(2): 228-232.

WILSON, M. (1997). Biocontrol of aerial plant diseases in agriculture and horticulture: current approaches and future prospects. *Journal of Industrial Microbiology and Biotechnology*, 19: 188–191.

YANG, D.Q., PLANTE, F., BERNIER, L., PICHÉ, Y., DESSUREAULT, M., LAFLAMME, G., OUELLETTE, G.B. (1993). Evaluation of a fungal antagonist, *Phaeothea dimorphospora*, for biological control of tree diseases. *Canadian Journal of Botany*, 71: 426-433.

YANG, D.Q., ROSSIGNOL, L. (1999). Evaluation of *Gliocladium roseum* against wood-degrading fungi in vitro and on major Canadian wood species. *Biocontrol Science and Technology*, 9: 409-420.

YANG, D.Q., WANG, X.M., WAN, H. (2007a). Biological protection of logs with *Gliocladium roseum* against biodegradation for panel manufacturing. *BioControl*, 52: 559-571.

YANG, D.Q., WANG, X.M., WAN, H. (2007b). Biological treatment of aspen strands to improve mold resistance and reduce resin consumption of strand boards. *Forest Products Journal*, 57(7/8): 58-62.

YANG, D.Q., WANG, X.M., WAN, H. (2007c). Use of fungal metabolites to protect wood based panels against mold infection. *BioControl*, 52: 427-436

YANG, D.Q. (2009). Biological Treatment to Improve Wood Product Quality and Durability – Fifteen Years of Effort and Experience at FPInnovations-Forintek Division. International Research Group on Wood Preservation, Document No. IRG/WP 09-40444.